


Steelhead smolt survival during the downstream and early ocean migration: effects of body size and migration distance

Michael Melnychuk, David Welch, Carl Walters



UBC




Marine mortality in steelhead trout

- Mortality rates during downstream and early ocean migration are probably high
- Growth vs. predation risk trade-offs
- Main groups of hypotheses for declines of total marine survival rates:
 - Reduced growth rates
 - Increased predation intensity






- Deployed array of VR2 receivers to cover >400 km of initial migratory routes
- Surgically implanted coded acoustic tags (69 kHz) into smolts during downstream migration
- Mobile tracking with VR-28 receiver/VFIN used on smaller spatial scale






- Deployed array of VR2 receivers to cover >400 km of initial migratory routes
- Surgically implanted coded acoustic tags (69 kHz) into smolts during downstream migration
- Mobile tracking with VR-28 receiver/VFIN used on smaller spatial scale



Howe Sound

2004
2005




Locations where tags
stopped moving:

Steelhead smolts

Howe Sound

2004

2005




Locations where tags
stopped moving:

Coho smolts


Results: survival rate estimation

- Mortality by population
- Body size effects
 - between species
 - between steelhead populations
 - within steelhead populations
- Migration distance effects
 - between steelhead populations



2004 %mortality
2005 %mortality

(wild)
FW: 81%



2004 %mortality

2005 %mortality

(wild)

FW: 81%

FW+O: 95%

Deadman
River

(wild)

FW: 31% , 14%

FW+O: 73% , 73%

Cheakamus
River

Coldwater
River

Keogh
River

(hatchery)

FW: 52% , 18%

FW+O: 71% , 42%

(wild)

FW: 48% , 58%

FW+O: 81% , 96%

(wild)

FW: 49%


FW+O: 69%


Englishman
River


(wild)


FW: 21% , 9%


FW+O: 85% , 88%












Cheakamus steelhead
(wild)




Steelhead

Mortality rate

1.0
0.8
0.6
0.4
0.2
0.0


12 14 16 18 20 22 24

Body length (cm)





FW

Steelhead




Steelhead








Downstream migration



Downstream + early ocean migration







51

Tagging and release point

Cheakamus River 3

Squamish River 4

Squamish River 5


Squamish River 6

Inner Howe

Sound line
(HS in)Outer Howe
Sound lines
(HS out)


10 km


34



Mortality rate within segment

● 2004
● 2005





2004 %mortality

2005 %mortality

(wild)

FW: 81%

FW+O: 95%

Deadman
River

(wild)

FW: 31% , 14%

FW+O: 73% , 73%

Cheakamus
River

Coldwater
River

Keogh
River

(hatchery)

FW: 52% , 18%

FW+O: 71% , 42%

(wild)

FW: 49%

FW+O: 69%

Englishman
River

(wild)

FW: 21% , 9%

FW+O: 85% , 88%

(wild)

FW: 48% , 58%

FW+O: 81% , 96%

2004 %mortality

2005 %mortality

(wild)

FW: 81%

FW+O: 95%

Deadman
River

(wild)

FW: 31% , 14%

FW+O: 73% , 73%

Cheakamus
River

Coldwater
River

Keogh
River

(hatchery)

FW: 52% , 18%

FW+O: 71% , 42%

(wild)

FW: 49%

FW+O: 69%

Englishman
River

(wild)


FW: 21% , 9%

FW+O: 85% , 88%

(wild)

FW: 48% , 58%

FW+O: 81% , 96%



**Sonia Batten
Isabelle Gaboury
Melinda Jacobs
Erin Rechisky**

**Jayson Semmens
Adrian Ladouceur
Paul Winchell
Caroline Melville**

Villy Christensen & UBC volunteers for mobile tracking

**Equipment: Fred Goetz (USACE), Martin Castonguay (DFO),
Gilles Lacroix (DFO)**

**Funding: Alfred P. Sloan Foundation (Census of Marine Life),
Gordon and Betty Moore Foundation, BPA,
FishAmerica Foundation, Pacific Salmon Foundation,
NSERC**

Thank you !

mikem@zoology.ubc.ca

